

A NEW VARIETY OF *PERYMENTIUM HINTONIORUM* (ASTERACEAE,
HELIANTHEAE)

B.L. Turner

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

A new variety of *Perymenium hintoniorum*, *P. h.* var. *gypsophilum* B.L. Turner, is described from southern Nuevo León. It is seemingly confined to gypsum soils and differs from the more northern var. *hintoniorum* of calcareous soils in having eglandular peduncles, smaller leaves and brown anthers. A map showing their distributions is provided.

KEY WORDS: Asteraceae, Heliantheae, *Perymenium*, México, Nuevo León, systematics

Routine identification of Mexican Asteraceae has revealed the following novelty.

PERYMENTIUM HINTONIORUM B.L. Turner var. ***GYPSOPHILUM*** B.L. Turner, var. nov. TYPE: MEXICO. Nuevo León: Mpio. Aramberri, along road from Aramberri to El Salitre, 1325 m, 26 Oct 1993, *Hinton et al.* 23749 (HOLOTYPE: TEX!).

A *P. hintoniorum* B.L. Turner var. *hintoniorum* folia parviora, 3-4 cm longa (vice 8-10 cm longa), et antheras brunneas (vice lotearum) habendo et solum en soliis gypseis (vice calcareorum) crescendo diagnosticum.

Suffruticose much-branched perennial herbs or shrublets 40-60 cm high. Leaves mostly 3-4(-7) cm long; petioles 2-8 mm long; blades ovate, having 3 principal veins, pubescent above and below with coarse hispid to pilose hairs, the margins serrate to nearly entire. Heads single on eglandular, sparsely strigose, peduncles 3-6 cm long. Involucres ca. 6 mm high, 6-10 mm wide (pressed); bracts 3-seriate, moderately strigose, the outer series broadly ovate, ca. 4 mm long, the inner lanceolate, ca. 6 mm long. Receptacle plane, the bracts linear-lanceolate, persistent. Ray florets 5-11, the ligules yellow. Disk florets 25-40; corollas yellow, ca. 6 mm long, glabrous except



Figure 1. Distribution of varieties of *Perymenium hintoniorum*.

for the hispidulous lobes. Anthers brown. Achenes 3-angled (ray florets) to radially flattened (disk florets), ca. 3 mm long, 1.8 mm wide; pappus of 10-20 deciduous bristles 2-4 mm long.

ADDITIONAL SPECIMENS EXAMINED: MEXICO. Nuevo León: Mpio. Aramberri, S of La Escondida, gypsum hillside, 1820 m, 16 Oct 1993, *Hinton et al.* 23596 (TEX); Aramberri to El Salitre, gypsum hillside, 1325 m, 26 Oct 1993, *Hinton et al.* 23733 (TEX); road to Dolores (from Aramberri), gypsum hills, 1255 m, *Hinton et al.* 23855 (TEX). Mpio. Zaragoza, W of Zaragoza, gypsum hillside, 1460 m, 16 Oct 1993, *Hinton et al.* 23645 (TEX); 19 km S of Zaragoza ($23^{\circ} 40' N$, $99^{\circ} 48' W$), 1950 m, 18 Nov 1993, *Villarreal y Carranza* 7560 (TEX).

As indicated in the diagnosis, var. *gypsophilum* is distinguished from the typical variety in having smaller leaves, eglandular peduncles and brown anthers. So far as known, it is confined to gypseous soils of southern Nuevo León (Figure 1).

ACKNOWLEDGMENTS

I am grateful to Gayle Turner for the Latin diagnosis, and to her and Justin Williams for reviewing the paper.